

A classification

of hemolytic uremic syndrome and thrombotic thrombocytopenic purpura and related disorders

N Besbas¹, D Karpman², D Landau³, C Loirat⁴, W Proesmans⁵, G Remuzzi^{6,11}, G Rizzoni⁷, CM Taylor⁸, N Van de Kar⁹ and LB Zimmerhackl¹⁰ from the European Paediatric Research Group for HUS

¹Hacettep University, Ankara, Turkey; ²Lund University, Lund, Sweden; ³Soroka Medical Centre, Beer Sheva, Israel; ⁴Hôpital Robert Debré, Paris, France; ⁵University Hospital, Leuven, Belgium; ⁶Mario Negri Institute for Pharmacological Research, Bergamo, Italy; ⁷Ospedale Bambina Gesù, Rome, Italy; ⁸Birmingham Children's Hospital, Birmingham, UK; ⁹UMC St Radboud, Nijmegen, The Netherlands and ¹⁰University of Innsbruck, Innsbruck, Austria

The diagnostic terms hemolytic uremic syndrome (HUS) and thrombotic thrombocytopenic purpura (TTP) are based on historical and overlapping clinical descriptions. Advances in understanding some of the causes of the syndrome now permit many patients to be classified according to etiology. The increased precision of a diagnosis based on causation is important for considering logical approaches to treatment and prognosis. It is also essential for research. We propose a classification that accommodates both a current understanding of causation (level 1) and clinical association in cases for whom cause of disease is unclear (level 2). We tested the classification in a pediatric disease registry of HUS. The revised classification is a stimulus to comprehensive investigation of all cases of HUS and TTP and is expected to increase the proportion of cases in whom a level 1 etiological diagnosis is confirmed.

Kidney International (2006) **70**, 423–431. doi:10.1038/sj.ki.5001581; published online 14 June 2006

KEYWORDS: hemolytic uremic syndrome; thrombotic thrombocytopenic purpura; etiology

New concepts of causation suggest an improved diagnostic classification of hemolytic uremic syndrome (HUS)¹ and thrombotic thrombocytopenic purpura (TTP),² two clinically defined syndromes with the shared pathology of thrombotic microangiopathy (TMA).^{3,4} Broadly, subgroups of HUS/TTP can be diagnosed at two levels. The first is by etiology, the second by clinical features or associations. The proposed classification is therefore split into two sections: part 1 in which concepts of etiology are reasonably advanced, and part 2 where clinical associations are used to describe patients for whom the cause remains unclear (Table 1). The division is admittedly arbitrary, and advancing knowledge is expected to redistribute cases from part 2 to 1. Future revisions will be needed. Subgroups re-defined by etiology often have a subgroup-specific prognosis and require specific therapies.

PROPOSED CLASSIFICATION: PART 1, ETIOLOGY ADVANCED Shiga and verocytotoxin (shiga-like toxin)-producing bacteria

The major cause of HUS in childhood is infection with verocytotoxin (shiga-like toxin)-producing bacteria, usually enterohemorrhagic *Escherichia coli* (VTEC/STEC),⁵ and in some tropical regions *Shigella dysenteriae* type I.^{6,7} Verocytotoxin-producing *Citrobacter freundii*, has also been reported.⁸ In America and the UK most cases are associated with *E. coli* serotype O157:H7,^{9–12} while other serotypes such as O26, O111, O103, and O145 are increasingly reported in Europe and elsewhere.^{13–15} The epidemiology has been extensively studied. Patients are usually pre-school children.^{16–18} VTEC strains produce various toxins, the major ones being verocytotoxin-1 (Stx1) and verocytotoxin-2 (Stx2). Verocytotoxin-1 differs by one amino acid from Shiga toxin produced by *Shigella dysenteriae* type 1. Verocytotoxin-2 has multiple variants that are closely related to each other, but have 55–60% homology to verocytotoxin-1. HUS is mostly caused by verocytotoxin-2-producing strains.^{11,14,15}

The disease begins after an incubation of 4–7 days with abrupt onset of diarrhea, usually bloody, with abdominal

Correspondence: CM Taylor, Department of Nephrology, Birmingham Children's Hospital, Birmingham B4 6NH, UK. E-mail: cm.taylor@bch.nhs.uk

¹¹Consultant to the European Paediatric Research Group for HUS.

Received 24 January 2006; revised 3 April 2006; accepted 18 April 2006; published online 14 June 2006

Table 1 | Classification of HUS, TTP, and related disorders

Part 1: etiology advanced	
1.i	<i>Infection induced</i> (a) Shiga and verocytotoxin (shiga-like toxin)-producing bacteria; enterohemorrhagic <i>Escherichia coli</i> , <i>Shigella dysenteriae</i> type 1, <i>Citrobacter</i> (b) <i>Streptococcus pneumoniae</i> , neuraminidase, and T-antigen exposure
1.ii	<i>Disorders of complement regulation</i> , (a) Genetic disorders of complement regulation (b) Acquired disorders of complement regulation, for example anti-FH antibody
1.iii	<i>von Willebrand proteinase, ADAMTS13 deficiency</i> (a) Genetic disorders of ADAMTS13 (b) Acquired von Willebrand proteinase deficiency; autoimmune, drug induced
1.iv	<i>Defective cobalamine metabolism</i>
1.v	<i>Quinine induced</i>
Part 2: Clinical associations: etiology unknown	
2.i	HIV
2.ii	Malignancy, cancer chemotherapy and ionizing radiation
2.iii	Calcineurin inhibitors and transplantation
2.iv	Pregnancy, HELLP syndrome and oral contraceptive pill
2.v	Systemic lupus erythematosus and antiphospholipid antibody syndrome
2.vi	Glomerulopathy
2.vii	Familial, not included in part 1
2.viii	Unclassified

FH, factor H; HELLP, HEMolytic anemia, elevated Liver enzymes, and Low Platelets; HIV, human immunodeficiency virus; HUS, hemolytic uremic syndrome; TTP, thrombocytopenic purpura.

pain. Microangiopathic hemolytic anemia, thrombocytopenia and acute oliguric renal failure occur 2–10 days later. The diagnosis of VTEC infection is made on stool culture, identification of toxin in the stools, or by serological response to the relevant O-serotype. Verotoxin itself is implicated in the pathogenesis, both from the epidemiology and from laboratory models. A prothrombotic state evolves prior to the acute renal failure,¹⁹ and the specific pathological finding is glomerular thrombosis.^{20,21}

With supportive therapy the mortality in children is <5% (*Shigella dysenteriae* type 1 excepted). Approximately 75% appear to make a full recovery when assessed up to 5 years after onset.²² Relapse is extremely rare.

***Streptococcus pneumoniae*, neuraminidase, and T-antigen exposure**

This is a distinctive disorder complicating pneumococcal infection, usually septicaemia, pneumonia with empyema, or meningitis with subdural collections.²³ Patients, usually <2 years old, present with marked microangiopathic hemolytic anemia. The acute mortality is about 25%.¹⁶ HUS relapse has not been described.

Pneumococcal neuraminidase has been identified in plasma.²⁴ It causes de-sialation of the glycocalyx of many

cells exposing the Thomsen–Friedenreich (T-) antigen that is normally covered by sialic acid.²⁵ T-antigen exposure on red cells is detected using the lectin *Hypogaea*. An immunoglobulin M cold antibody occurring naturally in human serum causes polyagglutination *in vitro*. Unlike other forms of HUS there is a positive Coomb's test. T-anti-T interaction on red cells, platelets, and endothelium was thought to explain the pathogenesis, but the role of the anti-T cold antibody *in vivo* is questionable.²⁶

Genetic disorders of complement regulation

Mutations in the genes for complement factor H (FH), factor I (FI), and membrane co-factor protein (MCP), also known as CD46, are associated with HUS.

FH, which is abundant in plasma, binds to polyanionic sites on host cells surfaces. Here it captures C3b generated by the alternative pathway and prevents formation of the C3 convertase, C3bBb, by factor B. This prevents amplification of the complement cascade and is a major reason why host cell surfaces are normally defended from alternative pathway activity.²⁷ FH is known to bind to vascular endothelium, erythrocytes, and platelets.^{28–30}

FH mutations have been found in a fifth of families with HUS and in about 8% in sporadic cases.^{31,32} Most patients have heterozygous mutations that affect the C-terminal, reducing binding to C3b/C3d, heparin,³³ and endothelial cells.^{29,34,35} Mutations leading to truncation of the FH molecule or the inability to be exported from the endoplasmic reticulum have also been described.³⁶ An association between three frequent single nucleotide polymorphisms of FH and susceptibility to HUS has been suggested.^{37,38}

The phenotype is variable and HUS may present at any age, although homozygous or compound heterozygous FH-deficient patients are more likely to manifest HUS in early life.³⁹ The plasma concentration of C3 may be normal or persistently low. Striking features include severe hypertension, a high risk of relapse, and poor prognosis, 50% of cases progressing to end-stage renal failure. The risk of graft loss after transplantation approaches 80%, mostly reflecting disease recurrence.^{40,41}

Mutations have been found in the genes encoding both FI^{42,43} and MCP (CD46).^{44,45} FI, a co-factor for FH, cleaves C3b interrupting the complement cascade before the generation of the anaphylotoxin C5a and the membrane attack complex C5b–C9 (reviewed by Zipfel *et al.*⁴⁶). Complement C5a and the membrane attack complex play an essential pathogenetic role in certain laboratory models of HUS.^{47,48}

FI circulates in plasma and, using FH, MCP, or CR1 as a co-factor, cleaves C3b to iC3b, or, using C4-binding protein, MCP or CR1 as a co-factor, cleaves C4b. All HUS patients so far described except one⁴³ are heterozygous for stop codons that truncate protein production, and plasma concentrations of FI may be reduced. MCP, a membrane-bound regulator expressed in glomerular endothelium also acts as a co-factor for the cleavage of C3b and C4b by FI. Heterozygous and homozygous MCP mutations have been identified in more

than 15 cases. These are predicted to cause loss of C3b and C4b recognition or loss of the C-terminal of the protein. Reduced expression on nucleated cells has been shown in some patients. Compared to FH or FI mutations, those with MCP abnormalities appear less likely to have disease recurrence after transplantation, probably because the donor organ retains normal MCP expression.

The diagnosis requires genetic expertise. A low C3 cannot be relied on to indicate deficient function of a membrane-bound complement regulator, nor does a normal plasma concentration of FH or FI exclude the possibility of a functional defect. Conversely, low C3 is a positive indicator, and in families with HUS those with hypocomplementemia are at increased risk of developing the disorder.⁴⁹ The fact that three different regulator abnormalities have been found so far supports the concept that failure to regulate complement on host tissues is etiologically relevant. It raises the possibility that other regulatory defects, either individual or compound,⁵⁰ will be found.

Acquired disorders of complement regulation

Evidence of immunoglobulin G autoantibodies to complement FH has been reported in three children with relapsing HUS.⁵¹ The plasma concentration of FH, and the FH gene were normal. FH autoantibody was identified by enzyme-linked immunosorbent assay, and shown to interfere with FH binding to the C3bBb convertase. These children also had antinuclear antibody but no other autoantibodies. Complement C3 was low in two cases as was factor B indicating alternative pathway activation. This strongly suggests that an acquired defect in complement regulation can mimic the inherited pattern outlined above.

Genetic disorders of von Willebrand factor-cleaving protease (ADAMTS13)

The finding of unusually large multimeric forms of von Willebrand factor (vWF), similar in size to those secreted by cultured endothelial cells *in vitro*, in the plasma of patients with HUS/TTP and chronic relapsing TTP was first reported by Moake *et al.*^{52,53} These large multimers cause platelet agglutination at high shear stress. In 1997, Furlan *et al.*⁵⁴ identified severe deficiency (<7% of normal plasma activity) of a specific protease (vWFCP) that cleaves multimeric vWF protein in patients with chronic relapsing TTP. This was confirmed in a large group of patients with familial and non-familial TTP.^{55,56} The vWFCP deficiency was shown to be either constitutional or due to autoantibodies. Lower activity occurs in liver disease, cancers, chronic inflammatory and metabolic conditions, pregnancy and in the newborn, but not the extremely low levels associated with TTP.^{57,58}

vWFCP cleaves multimeric vWF released from the endothelium and platelets into monomeric subunits at position 842–843. It has been purified, the gene sequenced,^{59,60} and shown to be a member of the ADAMTS family of metalloproteases and designated ADAMTS13. ADAMTS is an acronym for ‘a disintegrin-like and

metalloprotease with thrombospondin type I repeats’. ADAMTS13 is encoded on chromosome 9q34 and patients with familial relapsing TTP have mutations throughout the gene.^{61–64} vWFCP is produced by hepatic stellate cells^{65–67} and has a plasma half-life of 2–3 days.⁶⁸ Several assays of vWFCP activity and vWFCP-inhibiting autoantibodies (see below) have been developed.⁶⁹ Although reliable and comparable, they are labor intensive and specialized.

Chronic relapsing TTP due to inherited vWFCP deficiency presents with thrombocytopenia and hemolytic anemia usually in the neonatal period.^{63,70} Later, recurrent hemolysis and thrombocytopenia recur at intervals usually every 3rd–4th week. About 50% of the patients with vWFCP activity below 5% of normal have their first attack before the age of 5 years. In the others the diagnosis is made in adulthood,⁷¹ and not all patients present with the complete diagnostic criteria of HUS/TTP. If diagnosed early, chronic relapsing TTP/HUS due to constitutional vWFCP deficiency can be reversed or prevented by infusion of fresh-frozen or virus inactivated plasma or at 2–3 weeks interval without concurrent plasmapheresis.

Acquired disorders of vWF-cleaving protease

Severely decreased vWFCP activity in patients with acute acquired, non-familial TTP is mostly due to inhibitory immunoglobulin G autoantibodies.^{55,56} These are found in 48–80% of adult patients and are transient or intermittent in the majority. This form of TTP/HUS is mainly seen in adults and is usually a single acute episode, although recurrences have been reported in 11–36% of patients.

Historically, patients with untreated TTP had a mortality approaching 90%, and plasma exchange using fresh-frozen plasma reduced this to 20%.⁷² Patients with vWFCP deficiency due to autoantibodies usually respond to intensive plasmapheresis but may require additional immunosuppression or even splenectomy. Rituximab, the monoclonal antibody against CD20 on B lymphocytes is also effective.^{73–75}

Some cases of acquired ADAMTS13 deficiency have been linked to the platelet inhibitors ticlopidine and clopidogrel. Ticlopidine-associated TTP^{76,77} occurs in one per 1600–5000 patients treated, usually within 1 month of starting. Besides the hemolytic anemia and thrombocytopenia, 75% of cases developed neurological manifestations, and 30% renal insufficiency. Plasmapheresis reduces the mortality.⁷⁷ Autoantibodies inhibiting ADAMTS13 have been detected in ticlopidine-associated TTP.⁷⁸ Clopidogrel-associated TTP usually presents within 2 weeks of starting the drug, is more prone to recurrence than with ticlopidine, and requires more plasma exchanges. The incidence of clopidogrel-associated TTP is similar to the general population (about 3.7 cases per million) making a causal relation less certain. However, antibody-induced ADAMTS13 deficiency has been reported in two cases.⁷⁹

Defective cobalamin metabolism

This autosomal recessive form of HUS is attributed to an inborn error of cobalamin-C metabolism.⁸⁰ The biochemical characteristics of cobalamin-C deficiency are hyperhomocysteinemia and methylmalonic aciduria. High levels of homocysteine might be responsible for the vascular manifestations.

Patients with cobalamin-C deficiency usually present in the early days and months of life with failure to thrive, poor feeding, and vomiting. Rapid deterioration occurs due to metabolic acidosis, gastrointestinal bleeding, hemolytic anemia, thrombocytopenia, severe respiratory and hepatic failure, and renal insufficiency.^{81,82} It is likely that some die undiagnosed. Besides the early fulminant course, a more protracted disease can manifest later in childhood and adolescence.^{83–85} Renal biopsy showed a chronic TMA. Serum homocysteine can be 10 times normal values and urinary methylmalonic acid markedly increased but correct with daily hydroxycobalamin administration.⁸⁵

Quinine

Quinine, whether as a medication or food additive, is associated with TTP/HUS.^{86,87} In a series of HUS/TTP patients 11% reported taking quinine compared to 6% taking other drugs.⁸⁸ In sensitized patients the typical clinical pattern is an abrupt onset of chills, myalgia, vomiting, and oliguria immediately after quinine exposure. Anemia is often mild. Patients can be shown to have antibodies that recognize different glycoprotein epitopes on platelets, red cells, and leukocytes.^{89,90} This interaction is quinine dependent, suggesting that a neoantigen is formed. In platelets, the antibody has been found to cross-react with glycoprotein IIb/IIIa and sometimes Ib/IX. The disorder is comparatively mild and remits if plasma exchange is started early enough and quinine avoided. ADAMTS13 plasma activity is typically normal.

PROPOSED CLASSIFICATION: PART 2, CLINICAL ASSOCIATIONS; ETIOLOGY UNKNOWN

Human immunodeficiency virus

Human immunodeficiency virus (HIV) positivity and acquired immune deficiency syndrome are risk factors for renal disease⁹¹ and HIV nephropathy can include HUS. HUS appears more prevalent in the HIV population⁹² and has a high mortality in advanced acquired immune deficiency syndrome. In one series it occurred in a third of HIV-infected adults and was the most frequent cause of acute or rapidly progressive renal failure.⁹³ HIV-associated HUS/TTP is rarer in childhood.⁹⁴ Exceptionally, HIV-positive patients develop other forms of HUS, for example one patient had complete deficiency of vWF-cleaving protease ADAMTS13 having developed an antiprotease autoantibody,⁹⁵ but in the majority alternative explanations of causation have not been sought. It is assumed that the virus directly affects the endothelium causing TMA as has been shown in a primate model.⁹⁶

Malignancy, cancer chemotherapy, and ionizing radiation

Disseminated carcinoma, usually gastric,⁹⁷ prostatic,⁹⁸ or colon,⁹⁹ has been associated with HUS/TTP. In the registry of the Lombardi Cancer Research Center, Georgetown University, Washington DC, 85 patients met the criteria of hemolysis, thrombocytopenia, and serum creatinine above 1.6 mg/dl.¹⁰⁰ Eighty nine percent of them had adenocarcinoma, 26% being gastric cancer. HUS/TTP developed before cancer was diagnosed in a third of patients giving credence to the primary association. However, all except one received mitomycin at some point and it is unclear in many cases whether the association is directly with malignancy or with drugs used in its treatment. The same argument can be raised with irradiation. Infectious complications of cancer such as cytomegalovirus infection¹⁰¹ and human herpes virus-6¹⁰² may also play a role.

Mitomycin, an antimitotic used mainly in gastric and breast cancer, is associated with HUS and histological features of TMA.¹⁰³ Some cases improve with steroids and plasma exchanges.¹⁰⁴ Cattell¹⁰⁵ was able to induce TMA by injecting mitomycin into the renal artery of rats, supporting a causative role. Gemcitabine, a related anticancer drug, used for pulmonary, pancreatic, and urothelial carcinoma, is also thought to be the cause of HUS/TTP.^{106,107}

Radiation nephropathy presents with hypertension proteinuria or renal impairment up to a year after exposure. Cases with superadded microangiopathic hemolytic anemia typical of HUS are rare. The renal cells most sensitive to ionizing radiation are glomerular endothelium and tubular epithelium. In radiation nephropathy, electron microscopy shows that glomerular endothelial cells are swollen, separated from the basement membrane by electron-lucent material or lost entirely. A late finding is fibrinoid necrosis and thrombosis of arteries and arterioles. This resembles TMA adding credence to the association. Modern concepts of pathogenesis have not been applied.

Calcineurin inhibitors and transplantation

HUS/TTP occurs after transplantation of liver,¹⁰⁸ kidney,¹⁰⁹ heart, kidney-pancreas,¹¹⁰ and bone marrow, and with both cyclosporin-A and tacrolimus immunosuppression. In a review of 188 kidney transplants Zarifian *et al.*¹¹¹ calculated that incidence of TMA associated with cyclosporin-A was 14%, and with tacrolimus between 1 and 5%. With tacrolimus the clinical severity is very variable and drug levels do not predict development of TMA.¹¹² Dose reduction or changing one calcineurin antagonist for another¹¹³ sometimes results in recovery and suggests a causative role. These drugs exert both direct and endothelin-1-mediated vasoconstriction that reduces renal plasma flow and perhaps lead to prothrombotic changes in endothelium. However, the pathophysiology is speculative. Plasma activity of ADAMTS13 has not been measured in this setting.

HUS/TTP can occur up to several months after hematopoietic stem cell (bone marrow) transplantation. In a review, Moake and Byrnes¹¹⁴ point out that conditioning regimes with total body radiation and cyclophosphamide, and the use

of cyclosporin for graft-versus-host disease prophylaxis, make it impossible to determine whether any of these had a pathogenic role. George *et al.*¹¹⁵ conclude that the association between bone marrow transplantation and HUS/TTP is rare and that all in their series had evidence of underlying sepsis. The pathogenesis has not been convincingly shown. ADAMTS13 activity has been found to be decreased in patients undergoing allogeneic bone marrow transplantation but not to the very low levels associated with TTP, and in some the activity is normal.

It is unclear whether calcineurin inhibitors should be avoided in transplanting HUS patients from diagnostic subgroups known to have a high risk of relapse. In one report cyclosporin-A and tacrolimus were not associated with HUS recurrence.⁴¹ Sirolimus is of doubtful benefit¹¹⁶ as HUS recurred in two children treated with sirolimus while avoiding calcineurin inhibitors.¹¹⁷

Pregnancy, HELLP syndrome, and oral contraceptives

TTP is more prevalent in women than in men, and occurs more often in childbearing years. An association with the use of the oral contraceptive pill remains speculative, but an association with pregnancy is clear.¹¹⁸ HUS/TTP may present at any time during pregnancy but mostly in the last trimester and about the time of delivery. It is sometimes difficult to distinguish it from pre-eclampsia. Cases have been described in which TTP recurs in subsequent, although not always consecutive pregnancies. Patients with previous HUS/TTP not associated with pregnancy may or may not relapse when pregnant.

Comprehensive investigation of the cause of pregnancy-associated HUS/TTP has seldom been undertaken. A modest reduction in ADAMTS13 activity is induced by estrogens and occurs normally in the last trimester of pregnancy, but not to the very low levels observed in TTP.¹¹⁹ However, a few cases of pregnancy-associated HUS/TTP have severely reduced protease activity, with or without an inhibitor and the presence of ultralarge vWF multimers in plasma.^{120–123} Equally there are cases in which the protease is normal. There are no reports of complement activation in this group.

HELLP syndrome (HEmolytic anemia, elevated Liver enzymes, and Low Platelets) is also a disorder of the last trimester or parturition, and patients may have features of pre-eclampsia. The blood film typically has evidence of microangiopathic hemolytic anemia with fragmented red blood cells. Von Willebrand protease activity is reduced more than is seen in normal pregnancy, but again not to the low levels seen in TTP and without ultralarge vWF multimers in plasma.¹¹⁹ Whether the moderately reduced ADAMTS13 activity plays any role in the pathogenesis is unclear. One might predict that pregnancy, and pre-eclampsia in particular, would exacerbate a prothrombotic state and add to the causation of HUS/TTP, perhaps providing a 'second hit.'

Systemic lupus erythematosus, antiphospholipid antibody syndrome

HUS or TTP, systemic lupus erythematosus (SLE) and antiphospholipid antibody syndrome may coexist in the same

patient. HUS/TTP has been reported in 2–3%¹²⁴ and 8.4%¹²⁵ of SLE patients, with over 50 cases to date. It may manifest at any age, either before or years after SLE has been diagnosed.¹²⁶ The etiology is unclear. Autoantibodies to ADAMTS13, platelets and the platelet glycoprotein CD36 have been described in patients with SLE and it has been postulated that these incite endothelial injury and trigger the release of ultralarge vWF multimers, culminating in TMA.¹²⁴ Immunosuppressive and cytotoxic drugs have been used, in conjunction with plasmapheresis, to suppress production and increase clearance of these antibodies, with most patients achieving remission of microangiopathic symptoms. Some respond well to plasmapheresis alone.¹²⁴

HUS/TTP is a rare complication of antiphospholipid antibody syndrome and sometimes its presenting manifestation.¹²⁷ The main clinical consequences are severe hypertension, variable degrees of proteinuria, renal impairment, and cortical atrophy. Interestingly, in two patients severe ADAMTS13 deficiency was found due to autoantibodies.¹²⁸ Antiphospholipid antibodies have been suggested to increase the risk of TMA in SLE as lupus anticoagulant or anticardiolipin antibodies are found in the majority of SLE patients who develop HUS/TTP.¹²⁷

Glomerular capillary thrombosis is an additional vascular lesion that occurs in SLE, mainly in patients with antiphospholipid antibodies, and leads to glomerular sclerosis and renal insufficiency.¹²⁹ Steroids and plasma exchange alone or in combination, are the most common treatment. A meta-analysis showed that where steroids alone were the first treatment used, the clinical status and laboratory abnormalities worsened, whereas recovery occurred in 73% of episodes treated with plasma exchange.¹²⁷

Glomerular disorders

A small number of cases develop HUS superimposed on different forms of glomerular diseases. These are often children who are nephrotic at presentation^{130–134} although adults are also described.¹³⁵ HUS and membranoproliferative glomerulonephritis may coincide. Both are associated with alternative pathway complement activation and in a few cases where mutations affecting complement FH have been identified the classification would be better described as described in the section on Genetic disorders of complement regulation.

Familial/genetic disorders not included: Part 1

There are current familial cases extensively investigated for the causes advanced in part 1 of the proposed classification with negative results, indicating that other inherited risk factors await recognition.

Unclassified

Presently no cause or clinical association is found in the majority of childhood patients with non-diarrheal HUS, or in adults with HUS/TTP. Mostly these cases have not been comprehensively investigated. However, an undetermined proportion remains unclassified after full evaluation of complement regulation and ADAMTS13 activity.




Figure 1 | Sub-group diagnosis in 167 children with non-diarrheal, VTEC negative HUS entered into the registry of the European Pediatric Research Group. Part 1 subgroups are colored sections, part 2 are gray. Of the 34 with mutations in complement regulators, 22 involved FH, six FI, and six MCP. Two further cases had acquired anti-FH antibodies. Abnormality in ADAMTS13 protease and/or gene was found in 12 cases, two of whom also had a complement FH mutation. HUS attributable to pneumococcal sepsis occurred in 12 children. There were eight cases, four sibling pairs, in whom the etiology was unknown but the disorder was clearly familial. In one case HUS complicated existing mesangioproliferative glomerulonephritis. In 22 of the unclassified cases complement dysregulation was suspected because of relapsing disease and low C3 concentration, or abnormal Western blot profile of complement FH.

TESTING THE PROPOSED CLASSIFICATION

Figure 1 shows the diagnoses in 167 children with HUS in the European Paediatric Registry. Cases with diarrhea or VTEC infection (95% of all pediatric HUS) are excluded. In summary, 58 cases could be allocated to a level 1 diagnostic category, 35% of the total. Few of the 90 cases with no known causation have had complete investigation for complement dysregulation or ADAMTS13 activity.

DISCUSSION

Diagnostic terms need precision to be useful. HUS and TTP describe clinical presentations in the *absence* of knowledge about the cause, and even then may confuse. For example, there is no advantage in relabelling a case of VTEC-induced 'HUS' as 'TTP' because of neurological events that may affect over 10% of children. 'Atypical HUS,' often used to indicate presentation without diarrhea, is clearly a misnomer if it refers to a *typical* example of a known subgroup, for example HUS induced by pneumococcal sepsis. 'Intrinsic' and 'extrinsic' HUS/TTP is a laudable attempt to separate patients with host factors that promote recurrence and poor prognosis from those whose disease is the result of a single environmental insult.¹³⁶ However, this bipolar concept does not address any interplay between genetic and environmental factors. TMA is a pathological term and should be confined to this context.

Can the term TTP be adapted to indicate etiology and pathogenesis? Given that severe deficiency of ADAMTS13, has been identified mostly in those described clinically as TTP, while normal activity is usually found in VTEC-induced-, quinine-induced-, and bone marrow transplant-

associated HUS/TTP, this has been proposed. However, separation of TTP from HUS on the primary role of ADAMTS13 activity is not without its critics.¹³⁷ A wide range of ADAMTS13 activity can be found in various physiological and disease states, and a profound reduction has been shown in sepsis.¹³⁸ In an unselected series of 111 adult cases, severe ADAMTS13 deficiency provided 89% sensitivity and 91% specificity for what was clinically diagnosed as TTP.¹²² In another unselected series of 142 adults, only 18 cases had profound deficiency, and this finding did not distinguish them from the 32 with 'idiopathic HUS/TTP' without decreased activity.¹²³ Our proposal allows etiological and clinical descriptions to operate independently.

Some patients have two different etiologies, for example both VTEC-induced HUS and loss of ADAMTS13 activity,¹³⁹ decreased ADAMTS13 plus a FH mutation¹⁴⁰ and a FH mutation plus cobalamin deficiency.¹⁴¹ For any individual it may be necessary but not in itself sufficient that there is a mutation in FH, or a VTEC infection. There are good reasons to consider 'two-hit' or catastrophic models for disease expression. A mutated complement regulator may be a predisposition, VTEC infection a trigger. The proposed classification accommodates this way of thinking about causation. Dual causation will be missed if clinicians fail to explore further after a seemingly satisfactory diagnosis has been reached.

In any case of HUS/TTP, the history and the clinical pattern is important and may indicate an appropriate subgroup and therapy. Nevertheless, we propose that there should be comprehensive investigation of all cases with the only exception of a first episode of VTEC-positive, diarrhea-associated HUS in a child in whom there is no family history of HUS or TTP.

The difficulty in reaching an etiological diagnosis is considerable. Relevant tests have hitherto been available only through research laboratories. There is a pressing need for laboratory services to be available for the assessment of ADAMTS13 activity, and the genetic and functional investigation of complement regulation. To assist clinicians the European Paediatric Research Group for HUS maintains a website to indicate those laboratories that are able to assist (<http://espn.uwcm.ac.uk/guidelines.htm>).

An agreed classification is necessary if disease registries are to make progress in correlating clinical and etiological aspects. Without this, advice about treatment often lacks evidence,¹⁴² and treatment trials are difficult to design. We consider that the present classification will assist with this process. However, as no etiological diagnosis can be found in some thoroughly investigated cases, it is likely that additional diagnostic subgroups are still out there to be discovered. Our proposal is therefore an interim one destined for further development.

REFERENCES

1. Gasser C, Gautier E, Steck A et al. Hämolytisch-urämisches syndrome: bilaterale Nierenrinden nekrosen bei akuten erworbenen hämolytischen Anämien. *Schweiz Med Wochenschr* 1955; **85**: 905–909.

2. Moschcowitz E. An acute febrile pleiochromic anemia with hyalin thrombosis of the terminal arterioles and capillaries. An undescribed disease. *Arch Intern Med* 1925; **36**: 89-93.
3. Symmers WStC. Thrombotic microangiopathic haemolytic anaemia (thrombotic microangiopathy). *Br Med J* 1952; **2**: 897-903.
4. Habib R. Pathology of the hemolytic uremic syndrome. In: Kaplan B, Trompeter R, Moake J (eds). *Hemolytic Uremic Syndrome and Thrombotic Thrombocytopenic Purpura*. Marcel Dekker: New York, 1992, pp 315-353.
5. Karmali MA, Petric M, Lim C *et al*. The association between idiopathic hemolytic uremic syndrome and infection by verotoxin-producing *Escherichia coli*. *J Infect Dis* 1985; **151**: 775-782.
6. Butler T, Islam MR, Azad MA *et al*. Risk factors for development of hemolytic uremic syndrome during shigellosis. *J Pediatr* 1987; **110**: 894-897.
7. Bhimma R, Rollins NC, Coovadia HM, Adhikari M. Post-dysenteric hemolytic uremic syndrome in children during an epidemic of *Shigella* dysentery in Kwazulu/Natal. *Pediatr Nephrol* 1997; **11**: 560-564.
8. Tschape H, Prager R, Strekel W *et al*. Verotoxigenic *Citrobacter freundii* associated with severe gastroenteritis and cases of hemolytic uremic syndrome in a nursery school: green butter as the infection source. *Epidemiol Infect* 1995; **114**: 441-450.
9. Griffin P, Tauxe R. The epidemiology of infections caused by *Escherichia coli* O157:H7, other enterohemorrhagic *E. coli*, and the associated hemolytic uremic syndrome. *Epidemiol Rev* 1991; **13**: 60-98.
10. Rowe PC, Orrbine E, Wells GA *et al*. Epidemiology of hemolytic-uremic syndrome in Canadian children from 1986 to 1988. The Canadian Pediatric Kidney Disease Reference Centre. *J Pediatr* 1991; **119**: 218-224.
11. Milford D, Taylor CM, Guttridge B *et al*. Haemolytic uraemic syndromes in the British Isles 1985-8: association with verocytotoxin producing *Escherichia coli*. Part 1: clinical and epidemiological aspects. *Arch Dis Child* 1990; **65**: 716-721.
12. Decludt B, Bouvet P, Mariani-Kurkdjian P *et al*. Haemolytic uraemic syndrome and Shiga toxin-producing *Escherichia coli* infection in children in France. The Societe de Nephrologie Pediatrique. *Epidemiol Infect* 2000; **124**: 215-220.
13. Elliot EJ, Robins-Browne RM, O'Loughlin EV *et al*. Nationwide study of haemolytic uraemic syndrome: clinical, microbiological and epidemiological features. *Arch Dis Child* 2001; **85**: 125-131.
14. Friedrich AW, Bielaszewska M, Zhang WL *et al*. *Escherichia coli* harboring Shiga toxin 2 gene variants: frequency and association with clinical symptoms. *J Infect Dis* 2002; **185**: 74-84.
15. Beutin L, Krause G, Zimmermann S *et al*. Characterization of shiga toxin-producing *Escherichia coli* strains isolated from human patients in Germany over a 3-year period. *J Clin Microbiol* 2004; **42**: 1099-1108.
16. Lynn R, O'Brien SJ, Taylor CM *et al*. Childhood hemolytic uremic syndrome, United Kingdom and Ireland. *Emerg Infect Dis* 2005; **11**: 590-596.
17. Proulx F, Sockett P. Prospective surveillance of Canadian children with the haemolytic uraemic syndrome. *Pediatr Nephrol* 2005; **20**: 786-790.
18. Lopez EL, Diaz M, Grinstein S *et al*. Hemolytic uremic syndrome and diarrhea in Argentine children: role of shiga-like toxins. *J Infect Dis* 1989; **160**: 469-475.
19. Chandler W, Jelacic S, Boster D *et al*. Prothrombotic coagulation abnormalities preceding the hemolytic-uremic syndrome. *N Engl J Med* 2002; **346**: 23-32.
20. Richardson S, Karmali M, Becker L *et al*. The histopathology of the hemolytic uremic syndrome associated with verocytotoxin-producing *Escherichia coli* infections. *Hum Pathol* 1988; **19**: 1102-1108.
21. Inward CD, Howie AJ, Fitzpatrick MM *et al*. Renal histopathology in fatal cases of diarrhoea-associated haemolytic uraemic syndrome. *Pediatr Nephrol* 1997; **11**: 556-559.
22. Garg AX, Suri RS, Barrowman N *et al*. Long-term renal prognosis of diarrhea-associate haemolytic uremic syndrome. A systemic review, meta-analysis and meta- regression. *JAMA* 2003; **290**: 1360-1370.
23. Brandt J, Wong C, Mihm S *et al*. Invasive pneumococcal disease and hemolytic uremic syndrome. *Pediatrics* 2002; **110**: 371-376.
24. de Loos F, Huijben KMLC, Van der Kar NCAJ *et al*. Hemolytic uremic syndrome attributable to *Streptococcus pneumoniae* infection: a novel cause of secondary protein N-glycan abnormalities. *Clin Chem* 2002; **48**: 781-785.
25. Klein PJ, Bulla M, Newman RA *et al*. Thomsen-Friedenreich antigen in haemolytic uraemic syndrome. *Lancet* 1977; **2**: 1024-1025.
26. Eder A, Manno C. Does red-cell T activation matter? *Br J Haematol* 2001; **114**: 25-30.
27. Pangburn MK. Host recognition and target differentiation by factor H, a regulator of the alternative pathway of complement. *Immunopharmacology* 2000; **49**: 149-151.
28. Hellwage J, Jokiranta TS, Friesen MA *et al*. Complement C3b/C3d and cell surface polyanions are recognised by overlapping binding sites on the most carboxyl-terminal domain of complement factor H. *J Immunol* 2002; **169**: 6935-6944.
29. Manuelian T, Hellwage J, Meri S *et al*. Mutations in factor H reduce binding affinity to C3b and heparin and surface attachment to endothelial cells in haemolytic uremic syndrome. *J Immunol* 2002; **169**: 6935-6944.
30. Vaziri-Sani F, Hellwage J, Zipfel PF *et al*. Factor H binds to washed human platelets. *J Thromb Haemost* 2005; **3**: 154-162.
31. Warwicker P, Goodship THJ, Donne RL *et al*. Genetic studies into inherited and sporadic hemolytic uremic syndrome. *Kidney Int* 1998; **53**: 836-844.
32. Rougier N, Kazatchkine MD, Rougier J-P *et al*. Human complement factor H deficiency associated with hemolytic uremic syndrome. *J Am Soc Nephrol* 1998; **9**: 2318-2326.
33. Pangburn MK. Localisation of the host recognition functions of complement factor H at the carboxyl terminal: implications for hemolytic uremic syndrome. *J Immunol* 2002; **169**: 4702-4706.
34. Sanchez-Corral P, Gonzales-Rubio C, Rodrigues de Cordoba S, Lopez-Trascasa M. Functional analysis in serum from atypical hemolytic uremic syndrome patients reveals impaired protection of host cells associated with mutations in factor H. *Mol Immunol* 2004; **41**: 81-84.
35. Sanchez-Corral P, Perez-Caballero D, Huarte O *et al*. Structural and functional characterization of factor H mutations associated with atypical hemolytic uremic syndrome. *Am J Hum Genet* 2002; **71**: 1285-1295.
36. Ault BH, Schmidt BZ, Fowler NL *et al*. Human factor H deficiency. Mutations in framework cysteine residues and block in H protein secretion and intracellular catabolism. *J Biol Chem* 1997; **272**: 25168-25175.
37. Caprioli J, Castelletti F, Buccioni S *et al*. Complement factor H mutations and gene polymorphisms in haemolytic uraemic syndrome: the C-257T, the A2089G and the G2881T polymorphisms are strongly associated with the disease. *Hum Mol Genet* 2003; **15**: 3385-3395.
38. Esparza-Gordillo J, Goicoechea de Jorge E, Buil A *et al*. Predisposition to atypical haemolytic uremic syndrome involves the concurrence of different susceptibility alleles in the regulators of complement activation gene cluster in 1q32. *Hum Mol Genet* 2005; **14**: 703-712.
39. Landau D, Shalev H, Levy-Finer G *et al*. Familial haemolytic uremic syndrome associated with complement factor H deficiency. *J Pediatr* 2001; **138**: 12-17.
40. Loirat C, Naudet P. The risk of recurrent haemolytic uremic syndrome after renal transplantation in children. *Pediatr Nephrol* 2003; **18**: 1095-1101.
41. Bresin W, Daina E, Noris M *et al*. for the International Registry of Recurrent and Familial HUS/TTP. Outcome or renal transplantation in patients with non-Shiga toxin-associated haemolytic uremic syndrome. Prognostic significance of genetic background. *Clin J Am Soc Nephrol* 2006; **1**: 88-99.
42. Fremaux-Bacchi V, Dragon-Durey MA, Blouin J *et al*. Complement factor I: the susceptibility gene for atypical haemolytic uraemic syndrome. *J Med Genet* 2004; **41**: e84.
43. Kavanagh DM, Kemp EJ, Mayland E *et al*. Mutations in complement factor I predispose to development of atypical haemolytic uremic syndrome. *J Am Soc Nephrol* 2005; **16**: 2150-2155.
44. Richards A, Kemp EJ, Liszewski MK *et al*. Mutations in human complement regulator, membrane cofactor protein (CD46), predisposes to development of familial hemolytic uremic syndrome. *Proc Natl Acad Sci USA* 2003; **100**: 12966-12971.
45. Noris M, Brioschi S, Caprioli J *et al*. International registry of recurrence and familial HUS/TTP. Familial hemolytic uremic syndrome and an MCP mutation. *Lancet* 2003; **362**: 1542-1547.
46. Zipfel PF, Jokiranta TS, Hellwage J *et al*. The factor H protein family. *Immunopharmacology* 1999; **42**: 53-60.
47. Kondo C, Mizuno M, Nishikawa K *et al*. The role of C5a in the development of thrombotic glomerulonephritis in rats. *Clin Exp Immunol* 2001; **124**: 323-329.
48. Nangaku M, Alpers CE, Pippin J *et al*. CD59 protects glomerular endothelial cells from immune-mediated thrombotic microangiopathy in rats. *J Am Soc Nephrol* 1998; **9**: 590-597.
49. Noris M, Ruggerenti P, Perna A *et al*. Hypocomplementemia discloses genetic predisposition to hemolytic uremic syndrome and thrombotic

- thrombocytopenic purpura: role of factor H abnormalities. *J Am Soc Nephrol* 1999; **10**: 281–293.
50. Fremeaux-Bacchi V, Kemp EJ, Goodship JA et al. The development of atypical HUS is influenced by susceptibility factors in factor H and membrane cofactor protein- evidence from two independent cohorts. *J Med Genet* 2005; **42**: 852–856.
 51. Dragon-Durey M-A, Loirat C, Cloarec S et al. Anti-factor H autoantibodies associated with atypical haemolytic uremic syndrome. *J Am Soc Nephrol* 2005; **16**: 555–563.
 52. Moake JL, Byrnes CK, Troll JH et al. Unusually large plasma factor VIII: von Willebrand factor multimers in chronic relapsing thrombotic thrombocytopenic purpura. *N Engl J Med* 1982; **307**: 1432–1435.
 53. Moake JL, Byrnes JJ, Troll JH et al. Abnormal VIII: von Willebrand factor pattern in the plasma of patients with the haemolytic uremic syndrome. *Blood* 1984; **64**: 592–598.
 54. Furlan M, Robles R, Solenthaler M et al. Deficient activity of von Willebrand factor-cleaving protease in chronic relapsing thrombotic thrombocytopenic purpura. *Blood* 1997; **89**: 3097–3103.
 55. Furlan MF, Robles R, Galbusera M et al. Von Willebrand factor-cleaving protease in thrombotic thrombocytopenic purpura and the hemolytic uremic syndrome. *N Engl J Med* 1998; **339**: 1578–1584.
 56. Tsai HM, Chun-Yet Lain E. Antibodies to von Willebrand factor-cleaving protease in acute thrombotic thrombocytopenic purpura. *N Engl J Med* 1998; **338**: 1585–1594.
 57. Moake JL. Thrombotic microangiopathies. *N Engl J Med* 2002; **347**: 589–600.
 58. Mannucci PM, Canciani MT, Forza I et al. Changes in health and disease of the metalloprotease that cleaves von Willebrand factor. *Blood* 2001; **98**: 2730–2735.
 59. Fujikawa K, Suzuki H, McMullen B, Chung D. Purification of human von Willebrand factor-cleaving protease and its identification as a new member of the metalloproteinase family. *Blood* 2001; **98**: 1662–1666.
 60. Gerritsen HE, Robles R, Lämmle B, Furlan MF. Partial amino acid sequence of [purified von Willebrand factor-cleaving protease]. *Blood* 2001; **98**: 1654–1661.
 61. Levy GC, Nicols WC, Lian EC et al. Mutations in a member of the ADAMTS gene family cause thrombotic thrombocytopenic purpura. *Nature* 2001; **413**: 488–494.
 62. Kokame K, Matsumoto M, Soejima K et al. Mutations and common polymorphisms in ADAMTS13 gene responsible for von Willebrand factor-cleaving protease activity. *Proc Natl Assoc Soc* 2002; **99**: 11902–11907.
 63. Assink K, Schiphorst R, Allford S et al. Mutation analysis and clinical implications of von Willebrand factor protease deficiency. *Kidney Int* 2003; **63**: 1995–1999.
 64. Schneppenheim R, Budde U, Oyen F et al. Von Willebrand factor cleaving protease and ADAMTS13 mutations in childhood TTP. *Blood* 2003; **101**: 1845–1850.
 65. Zheng X, Chung D, Takayama TK et al. Structure of von Willebrand factor-cleaving protease (ADAMTS13), a metalloprotease involved in thrombotic thrombocytopenic purpura. *J Biol Chem* 2001; **276**: 41059–41063.
 66. Soejima K, Mimura M, Hirashima M. A novel human metalloprotease synthesized in the liver and secreted into the blood: possibly, the von Willebrand factor-cleaving protease? *J Biochem* 2001; **130**: 475–480.
 67. Uemura M, Tatsumi K, Matsumoto M et al. Localization of ADAMTS13 to the stellate cells of human liver. *Blood* 2005; **106**: 922–924.
 68. Furlan M, Robles R, Morselli B et al. Recovery and half life of von Willebrand factor-cleaving-protease after plasmapheresis in patients with thrombotic thrombocytopenic purpura. *Thromb Haemost* 1999; **81**: 8–13.
 69. Tripodi A, Chantarangkul V, Bohm M et al. Measurements of von Willebrand factor cleaving protease (ADAMTS13): results of an international collaborative study involving 11 methods testing the same set of coded plasmas. *J Thromb Haemost* 2004; **2**: 1601–1609.
 70. Jubinsky PT, Moraille R, Tsai HM. Thrombotic thrombocytopenic purpura in a newborn. *J Perinatol* 2003; **23**: 85–87.
 71. George JN, Sadler JE, Lämmle B. Platelets: thrombotic thrombocytopenic purpura. *Hematology (Am Soc Hematol Educ Program)* 2002: 315–342.
 72. Rock GA, Shumak KH, Buskard NA et al. Comparison of plasma exchange with plasma infusion in the treatment of thrombotic thrombocytopenic purpura. A Canadian apheresis study group. *N Engl J Med* 1991; **325**: 393–397.
 73. Zheng X, Pallera AM, Goodnough LT et al. Remission of chronic thrombotic thrombocytopenic purpura after treatment with cyclophosphamide and rituximab. *Ann Intern Med* 2003; **138**: 105–108.
 74. Tsai HM. Advances in the pathogenesis, diagnosis, and treatment of thrombotic thrombocytopenic purpura. *J Am Soc Nephrol* 2003; **14**: 1072–1081.
 75. Fakhouri F, Vernant JP, Veyradier A et al. Efficiency of curative and prophylactic treatment with rituximab in ADAMTS13-deficient thrombotic thrombocytopenic purpura: a study of 11 cases. *Blood* 2005; **106**: 1932–1937.
 76. Bennett CL, Weinberg PD, Rozenberg-Ben-Dror K et al. Thrombotic thrombocytopenic purpura associated with ticlopidine. A review of 60 cases. *Ann Intern Med* 1998; **128**: 541–544.
 77. Bennett CL, Davidson CJ, Risch DW et al. Thrombotic thrombocytopenic purpura associated with ticlopidine in the setting of coronary artery stents and stroke prevention. *Ann Intern Med* 1999; **159**: 2524–2528.
 78. Tsai HM, Rice L, Sarode R et al. Antibody inhibitors to von Willebrand factor metalloproteinase and increased binding of von Willebrand factor to platelets in ticlopidine-associated thrombotic thrombocytopenic purpura. *Ann Intern Med* 2000; **132**: 7984–7989.
 79. Bennett CL, Connors JM, Carwile JM et al. Thrombotic thrombocytopenic purpura associated with clopidogrel. *N Engl J Med* 2000; **342**: 1773–1777.
 80. Baumgartner ER, Wick H, Maurer R et al. Congenital defect in intracellular cobalamin metabolism resulting in homocystinuria and methylmalonic aciduria. A case report and histopathology. *Helv Paediatr Acta* 1979; **34**: 465–482.
 81. Russo PA, Doyon J, Sonsino E et al. Inborn errors of cobalamin metabolism and the hemolytic uremic syndrome. In: Kaplan BS, Trompeter RS, Moake JL (eds). *Hemolytic Uremic Syndrome and Thrombotic Thrombocytopenic Purpura*. Dekker: New York, 1992, pp 255–270.
 82. Geraghty MT, Perlman EJ, Martin LS et al. Cobalamin C defect associated with hemolytic-uremic syndrome. *J Pediatr* 1992; **120**: 934–937.
 83. Labrune P, Zittoun J, Duvalier I et al. Haemolytic uraemic syndrome and pulmonary hypertension in a patient with methionine synthase deficiency. *Eur J Pediatr* 1999; **158**: 734–739.
 84. Brunelli SM, Meyers KEC, Guttenberg M et al. Cobalamin C deficiency complicated by an atypical glomerulopathy. *Pediatr Nephrol* 2002; **17**: 800–803.
 85. Van Hove JL, Van Damme-Lombaerts R, Grunewald S et al. Cobalamin disorder Cbl-C presenting with late-onset thrombotic microangiopathy. *Am J Med Genet* 2002; **111**: 195–201.
 86. Gottsall JL, Neahring B, MacFarland JG et al. Quinine-induced immune thrombocytopenia with haemolytic uremic syndrome: clinical and serological findings in nine patients and review of literature. *Am J Hematol* 1994; **47**: 283–289.
 87. Aster RH. Quinine sensitivity. A new cause of the haemolytic uremic syndrome. *Ann Int Med* 1993; **119**: 243–244.
 88. Kojouri K, Vesely SK, George JN. Quinine-associated thrombotic thrombocytopenic purpura-hemolytic uremic syndrome: frequency, clinical features and long-term outcomes. *Ann Intern Med* 2001; **135**: 1047–1051.
 89. Stroncek DF, Vercellotti GM, Hammerschmidt DE et al. Characterization of multiple quinine-dependent antibodies in a patient with episodic haemolytic uremic syndrome and immune agranulocytosis. *Blood* 1992; **80**: 241–248.
 90. Glynne P, Salama A, Chaudhry A et al. Quinine-induced immune thrombocytopenic purpura followed by hemolytic uremic syndrome. *Am J Kidney Dis* 1999; **33**: 133–137.
 91. Strauss J, Abitbol C, Zilleruelo G, Montane B. HIV nephropathy. In: Barratt TM, Avner ED, Harmon WE (eds). *Pediatric Nephrology*. 4th edn, Lippincott, Williams and Wilkins: Baltimore, 1999, pp 1103–1107.
 92. Badesha PS, Saklayen MG. Hemolytic uremic syndrome as a presenting form of HIV infection. *Nephron* 1996; **72**: 472–475.
 93. Peraldi MN, Maslo C, Akposso K et al. Acute renal failure in the course of HIV infection: a single institution retrospective study of ninety-two patients and sixty renal biopsies. *Nephrol Dial Transplant* 1999; **14**: 1578–1585.
 94. Ray PE, Rakusan T, Loehelt BJ et al. Human immunodeficiency virus (HIV) associated nephropathy in children from the Washington DC. Area: 12 years' experience. *Semin Nephrol* 1998; **18**: 396–405.
 95. Sahud MA, Claster S, Liu L et al. Von Willebrand factor cleaving protease inhibitor in a patients with human immunodeficiency syndrome

- associated thrombotic thrombocytopenic purpura. *Br J Haematol* 2002; **116**: 909–911.
96. Segerer S, Eitner F, Cui Y et al. Cellular injury associated with renal thrombotic microangiopathy in human immunodeficiency virus-infected macaques. *J Am Soc Nephrol* 2002; **13**: 370–378.
 97. Carr DJ, Kramer BS, Dragonetti DE. Thrombotic thrombocytopenic purpura associated with metastatic gastric adenocarcinoma: successful management with plasmapheresis. *South Med J* 1986; **79**: 476–479.
 98. Mungall S, Mathieson P. Hemolytic uremic syndrome in metastatic adenocarcinoma of the prostate. *Am J Kidney Dis* 2002; **40**: 1334–1336.
 99. Majhail NS, Hix JK, Almahameed A. Carcinoma of the colon in a patient presenting with thrombotic thrombocytopenic purpura-hemolytic uremic syndrome. *Mayo Clin Proc* 2002; **77**: 873.
 100. Lesesne JB, Rothschild N, Erickson B et al. Cancer-associated hemolytic-uremic syndrome: analysis of 85 cases from a national registry. *J Clin Oncol* 1989; **7**: 781–789.
 101. Cavagnaro F, Barriga F. Hemolytic uremic syndrome in a child with leukemia and cytomegalovirus infection. *Pediatr Nephrol* 2000; **14**: 1118–1120.
 102. Matsuda Y, Hara J, Miyoshi H et al. Thrombotic microangiopathy associated with reactivation of human herpesvirus-6 following high-dose chemotherapy with autologous bone marrow transplantation in young children. *Bone Marrow Transplant* 1999; **25**: 919–923.
 103. Crocker J, Jones EL. Haemolytic-uraemic syndrome complicating long-term mitomycin C and 5-fluorouracil therapy for gastric carcinoma. *J Clin Pathol* 1983; **36**: 24–29.
 104. Lyman NW, Michaelson R, Viscuso RL et al. Successful treatment with corticosteroids and intense plasma exchange. *Arch Intern Med* 1983; **143**: 1617–1618.
 105. Cattell V. Mitomycin-induced hemolytic uremic kidney. An experimental model in the rat. *Am J Pathol* 1985; **121**: 88–95.
 106. Teixeira L, Debourdeau P, Zammit C et al. Gemcitabine-induced thrombotic microangiopathy. *Presse Méd* 2002; **31**: 740–742.
 107. Humphreys BD, Sharman JP, Henderson JM et al. Gemcitabine-associated thrombotic microangiopathy. *Cancer* 2004; **100**: 2664–2670.
 108. Bonser RS, Adu D, Franklin I, McMaster P. Cyclosporin-induced haemolytic uraemic syndrome in liver allograft recipient. *Lancet* 1984; **2**: 1337.
 109. Van Buren D, Van Buren CT, Flechner SM et al. De novo hemolytic uremic syndrome in renal transplant recipients immunosuppressed with cyclosporine. *Surgery* 1985; **98**: 54–62.
 110. Young BA, Marsh CL, Alpers CE, Davis CL. Cyclosporine-associated thrombotic microangiopathy/hemolytic uremic syndrome following kidney and kidney-pancreas transplantation. *Am J Kidney Dis* 1996; **28**: 561–571.
 111. Zarifian A, Meleg-Smith S, O'donovan R et al. Cyclosporine-associated thrombotic microangiopathy in renal allografts. *Kidney Int* 1999; **55**: 2457–2466.
 112. Trimarchi HN, Truong LD, Brennan S et al. FK506-associated thrombotic microangiopathy. Report of two cases and review of the literature. *Transplantation* 1999; **67**: 539–544.
 113. Burke GAA, McGraw ME, MacIver AG. The role of early renal biopsy in cyclosporin induced thrombotic microangiopathy. *Pediatr Nephrol* 1999; **13**: 564–566.
 114. Moake JL, Byrnes JJ. Thrombotic microangiopathies associated with drugs and bone marrow transplantation. *Hematol Oncol Clin North Am* 1996; **10**: 485–497.
 115. George JN, Li X, McMin J et al. Thrombotic thrombocytopenic purpura – haemolytic uremic syndrome following allogeneic hematopoietic stem cell transplantation: a diagnostic dilemma. *Transfusion* 2004; **44**: 294–304.
 116. Reynolds JC, Agodoa LY, Yuam CM, Abbott KC. Thrombotic microangiopathy after renal transplantation in the United States. *Am J Kidney Dis* 2003; **42**: 1058–1068.
 117. Florman S, Benchimol C, Lieberman K et al. Fulminant recurrence of atypical haemolytic uremic syndrome during a calcineurin inhibitor-free immunosuppression regimen. *Pediatr Transpl* 2002; **6**: 352–355.
 118. George JN, Vesely SK, Terrell DR. The Oklahoma thrombotic thrombocytopenic purpura-hemolytic uremic syndrome (TTP-HUS) registry: a community perspective of patients with clinically diagnosed TTP-HUS. *Semin Hematol* 2004; **41**: 60–67.
 119. Lattuada A, Rossi E, Calzarossa R, Mannucci PM. Mild to moderate reduction of a von Willebrand factor cleaving protease (ADAMTS-13) in pregnant women with HELLP microangiopathic syndrome. *Haematologica* 2003; **88**: 1029–1034.
 120. Cosmai EM, Puzis L, Tsai H-M, Lian EC-Y. Thrombocytopenic purpura and cardiomyopathy in pregnancy reversed by combined plasma exchange and infusion. *Eur J Haematol* 2002; **68**: 239–242.
 121. Ducloy-Bouthors A-S, Caron C, Subtil D et al. Thrombotic thrombocytopenic purpura: medical and biological monitoring of six pregnancies. *Eur J Obstet Gynecol Reprod Biol* 2003; **111**: 146–152.
 122. Veyradier A, Obert B, Houlier A et al. Specific von Willebrand factor-cleaving protease in thrombotic microangiopathies: a study of 111 cases. *Blood* 2001; **98**: 1765–1771.
 123. Vesely SK, George JN, Lammle B et al. ADAMTS13 activity in thrombotic thrombocytopenic purpura-hemolytic uremic syndrome: relation to presenting features and clinical outcomes in a prospective cohort of 142 patients. *Blood* 2003; **102**: 60–68.
 124. Vasoo S, Thumboo J, Fong KY. Thrombotic thrombocytopenic purpura in systemic lupus erythematosus: disease activity and the use of cytotoxic drugs. *Lupus* 2002; **11**: 443–450.
 125. Banfi G, Bertani T, Boeri V et al. Renal vascular lesions as a marker of poor prognosis in patients with lupus nephritis. *Am J Kidney Dis* 1991; **18**: 240–248.
 126. Kawasaki Y, Suzuki J, Nozawa R et al. A 12 year-old girl with hemolytic uremic syndrome as initial symptom of systemic lupus erythematosus and a literature review. *Am J Nephrol* 2002; **22**: 576–580.
 127. Espinosa G, Bucciarelli S, Cervera R et al. Thrombotic microangiopathic haemolytic anaemia and antiphospholipid antibodies. *Ann Rheum Dis* 2004; **63**: 730–736.
 128. Amoura Z, Costedoat-Chalumeau N, Veyradier A et al. Thrombotic thrombocytopenic purpura with severe ADAMTS-13 deficiency in two patients with primary antiphospholipid syndrome. *Arthritis Rheum* 2004; **50**: 3260–3264.
 129. Ruggerenti P, Galli M, Remuzzi G. Hemolytic uremic syndrome, thrombotic thrombocytopenic purpura and antiphospholipid antibody syndromes. In: Neilson EG and Couser WG (eds). *Immunologic Renal Diseases*. Lippincott Williams and Wilkins: Philadelphia, 2001, Chapter 49, pp 1173–1201.
 130. De Chadarevian JP, Goodyer PR, Kaplan BS. Acute glomerulonephritis and hemolytic uremic syndrome. *Can Med Assoc J* 1980; **123**: 391–394.
 131. Proesmans W, Baten E, Van Damme B. A boy with acute renal failure. A case for diagnosis. *Pediatr Nephrol* 1995; **9**: 389–391.
 132. Habib R. Pathology of the hemolytic uremic syndrome. In: Kaplan BS, Trompeter RS and Moake JL (eds). *Hemolytic Uremic Syndrome and Thrombotic Thrombocytopenic Purpura*. Dekker: New York, 1992, pp 341–342.
 133. Siegler RL, Brewer ED, Pysher J. Hemolytic uremic syndrome associated with glomerular disease. *Am J Kidney Dis* 1989; **13**: 144–147.
 134. Dische FE, Culliford EJ, Parsons V. Haemolytic urmic syndrome and idiopathic membranous glomerulonephritis. *Br Med J* 1978; **1**: 1112–1113.
 135. Morita S, Sakai T, Okamoto N et al. Hemolytic uremic syndrome associated with immunoglobulin A nephropathy: a case report and review of cases of haemolytic uremic syndrome with glomerular disease. *Intern Med* 1999; **38**: 495–499.
 136. Melnyk AM, Solez K, Kjellstrand CM. Adult hemolytic-uremic syndrome. A review of 37 cases. *Arch Intern Med* 1995; **155**: 2077–2084.
 137. Remuzzi G, Galbusera M, Mannucci PM. ADAMTS13 in microangiopathies. *Blood* 2002; **100**: 3842.
 138. Ono T, Mimuro J, Madoiwa S et al. Severe deficiency of von Willebrand factor-cleaving protease (ADAMTS13) in patients with sepsis-induced disseminated intravascular coagulation: its correlation with development of renal failure. *Blood* 2006; **107**: 528–534.
 139. Hunt BJ, Lammle B, Nevard CHF, Haycock GB. Von Willebrand factor-cleaving protease in childhood diarrhoea-associated haemolytic uraemic syndrome. *Thromb Haemost* 2001; **85**: 975–978.
 140. Noris M, Bucchioni S, Galbusera M et al. Complement factor H mutation in familial thrombotic thrombocytopenic purpura with ADAMTS13 deficiency and renal involvement. *J Am Soc Nephrol* 2005; **16**: 1177–1183.
 141. Guignon V, Fremeaux-Bacchi V, Giraudier S et al. Late-onset thrombotic microangiopathy caused by cblC disease: association with a factor H mutation. *Am J Kidney Dis* 2005; **45**: 588–595.
 142. Allford SL, Hunt BJ, Rose P, Machin SJ. (on behalf of the haemostasis and thrombosis task force of the British Committee for Standards in Haematology) Guidelines on the diagnosis and management of the thrombotic microangiopathic haemolytic anaemias. *Br J Haematol* 2003; **120**: 556–573.